

Physics 566

Problem Set #6 Solutions

Problem 1: Boson Algebra

(a) $\int \frac{d^2\beta}{\pi} e^{-A|\beta|^2} e^{\alpha\beta^* - \alpha^*\beta}$ Let $\alpha = X_1 + iX_2$
 $\beta = Y_1 + iY_2$

$\int \frac{dY_1 dY_2}{\pi} e^{-A(Y_1^2 + Y_2^2)} e^{2i(X_2 Y_1 - X_1 Y_2)}$

$= \left(\int_{-\infty}^{\infty} \frac{dY_1}{\sqrt{\pi}} e^{-AY_1^2} e^{2iX_2 Y_1} \right) \left(\int_{-\infty}^{\infty} \frac{dY_2}{\sqrt{\pi}} e^{-AY_2^2} e^{2iX_1 Y_2} \right)$

(Gaussian Fourier Integral)

$\rightarrow \underbrace{e^{-X_2^2/A}}_{\frac{1}{\sqrt{A}}} \underbrace{e^{-X_1^2/A}}_{\frac{1}{\sqrt{A}}}$

$= \frac{1}{A} e^{-|\alpha|^2/A}$

(b) $\int \frac{d^2\alpha}{\pi} |\alpha\rangle\langle\alpha| = \sum_{n,m} \int \frac{d^2\alpha}{\pi} \left(\frac{\alpha^n}{\sqrt{n!}} e^{-|\alpha|^2/2} \right) \left(\frac{\alpha^{*m}}{\sqrt{m!}} e^{-|\alpha|^2/2} \right)$
 $|\alpha\rangle\langle m|$

change to polar coords: $\alpha = A e^{i\theta} \Rightarrow d^2\alpha = A d\theta dA$

$\Rightarrow \int \frac{d^2\alpha}{\pi} |\alpha\rangle\langle\alpha| = \sum_{n,m} \frac{|\alpha\rangle\langle m|}{\pi \sqrt{n!m!}} \int_0^\infty A dA e^{-A^2} A^{n+m} \int_0^{2\pi} d\theta e^{i(n-m)\theta}$
 $2\pi \delta_{nm}$

$= \sum_{n=0}^{\infty} \frac{2|\alpha\rangle\langle n|}{\pi n!} \int_0^\infty dA A^{2n+1} e^{-A^2}$
 $\frac{n!}{2}$ (Gamma Function)

$= \sum_n |\alpha\rangle\langle n| = \mathbb{1} \checkmark$

(c) Quadrature operators: $\hat{a} = \hat{X}_1 + i\hat{X}_2$

$$\Rightarrow \hat{X}_1 = (\hat{a} + \hat{a}^\dagger)/2 \quad \hat{X}_2 = (\hat{a} - \hat{a}^\dagger)/2i$$

Under $\hat{U} = e^{-i\theta \hat{a}^\dagger \hat{a}}$
 \uparrow
 free Hamiltonian

$$\hat{U}^\dagger \hat{a} \hat{U} = \hat{a} e^{-i\theta}$$

$$\hat{U}^\dagger \hat{a}^\dagger \hat{U} = \hat{a}^\dagger e^{+i\theta}$$

$$\begin{aligned} \Rightarrow \hat{U}^\dagger \hat{X}_1 \hat{U} &= \frac{1}{2} (\hat{U}^\dagger \hat{a} \hat{U} + \text{h.c.}) = \frac{1}{2} (\hat{a} e^{-i\theta} + \hat{a}^\dagger e^{i\theta}) \\ &= \text{Re}(\hat{a} e^{-i\theta}) = \hat{X}_1 \cos\theta + \hat{X}_2 \sin\theta \end{aligned}$$

$$\hat{U}^\dagger \hat{X}_2 \hat{U} = \text{Im}(\hat{a} e^{-i\theta}) = \hat{X}_2 \cos\theta - \hat{X}_1 \sin\theta$$

The free evolution operator corresponds to a rotation in phase space

(d) Consider $\hat{D}(\alpha + \beta) = \exp\{(\alpha + \beta)\hat{a}^\dagger - (\alpha^* + \beta^*)\hat{a}\}$

Ans: Recall Baker-Campbell-Hausdorff

$$e^{A+B} = e^A e^B e^{-\frac{1}{2}[A,B]} \quad \text{when } [A,B] \text{ commutes with } A \text{ and } B$$

$$\Rightarrow \hat{D}(\alpha + \beta) = \hat{D}(\alpha) \hat{D}(\beta) \exp\left\{-\frac{1}{2}[\alpha \hat{a}^\dagger - \alpha^* \hat{a}, \beta \hat{a}^\dagger - \beta^* \hat{a}]\right\}$$

$$\hat{D}(\alpha + \beta) = \hat{D}(\alpha) \hat{D}(\beta) \exp\left\{-\frac{1}{2}(\alpha \beta^* - \alpha^* \beta)\right\}$$

$i \text{Im}(\alpha \beta^*)$

$$\Rightarrow \boxed{\hat{D}(\alpha) \hat{D}(\beta) = \hat{D}(\alpha + \beta) \exp\{i \text{Im}(\alpha \beta^*)\}}$$

(e) Matrix elements of displacement operator

• Vacuum: $\langle 0 | \hat{D}(\alpha) | 0 \rangle = \langle 0 | e^{\alpha \hat{a}^\dagger - \alpha^* \hat{a}} | 0 \rangle$

$$= \underbrace{\langle 0 | e^{\alpha \hat{a}^\dagger} e^{-\alpha^* \hat{a}} | 0 \rangle}_{=1} e^{-|\alpha|^2/2} \quad (\text{Normal order})$$

• Coherent States: $\langle \alpha_1 | \hat{D}(\alpha) | \alpha_2 \rangle = \langle 0 | \hat{D}^\dagger(\alpha_1) \hat{D}(\alpha) \hat{D}(\alpha_2) | 0 \rangle$

$$= \langle 0 | \hat{D}^\dagger(-\alpha_1) \hat{D}(\alpha) \hat{D}(\alpha_2) | 0 \rangle = \langle 0 | \hat{D}(\alpha_1) \hat{D}(\alpha + \alpha_2) | 0 \rangle e^{i \text{Im}(\alpha \alpha_2^*)}$$

$$= \langle 0 | \hat{D}(\alpha + \alpha_2 - \alpha_1) | 0 \rangle \exp\{i \text{Im}(\alpha \alpha_2^*) - i \text{Im}(\alpha_1(\alpha^* + \alpha_2^*))\}$$

$$= \exp\left\{-\frac{|\alpha + \alpha_2 - \alpha_1|^2}{2}\right\} \exp\left\{i \text{Im}(\alpha \alpha_2^* - \alpha_1 \alpha^* - \alpha_1 \alpha_2^*)\right\}$$

• Fock states: $\langle n | \hat{D}(\alpha) | n \rangle = \langle n | e^{\alpha \hat{a}^\dagger} e^{-\alpha^* \hat{a}} | n \rangle e^{-|\alpha|^2/2}$

$$= \sum_{m, m'} \frac{(n!)^m}{m! m'} \frac{(\hat{a}^\dagger)^m (\hat{a})^{m'}}{m! m'} e^{\alpha^m} (\alpha)^{m'} e^{-|\alpha|^2/2}$$

$$= \underbrace{\left(\sum_{m=0}^{\infty} (-1)^m \frac{n!}{(n-m)!} \frac{1}{(m!)^2} (\alpha)^m \right)}_{\mathcal{L}_n(|\alpha|^2)} e^{-|\alpha|^2/2}$$

$\mathcal{L}_n(|\alpha|^2)$ see Appendix (B.32) 3rd edition

$$\Rightarrow \langle n | \hat{D}(\alpha) | n \rangle = e^{-|\alpha|^2/2} \mathcal{L}_n(|\alpha|^2)$$

Problem 2: Properties of the Wigner function

$$W(x_1, x_2) = \int \frac{dx'}{\pi} e^{-2ix'x_2} \langle x_1 - x' | \hat{\rho} | x_1 + x' \rangle$$

$$= \int \frac{dx'}{\pi} e^{-2ix'x_2} \psi(x_1 - x') \psi^*(x_1 + x')$$

The last form is what Wigner originally wrote down in considering the phase space description of ordinary quantum mechanics.

(a) Marginals: $P(x_1) = \int dx_2 W(x_1, x_2)$
 $P(x_2) = \int dx_1 W(x_1, x_2)$

We seek to show $P(x_1) = \langle x_1 | \hat{\rho} | x_1 \rangle = \left(|\psi(x_1)|^2 \right)$ Pure States
 $P(x_2) = \langle x_2 | \hat{\rho} | x_2 \rangle = \left(|\tilde{\psi}(x_2)|^2 \right)$ States

$$\int dx_2 W(x_1, x_2) = \int \frac{dx'}{\pi} \int e^{-2ix'x_2} dx_2 \langle x_1 - x' | \hat{\rho} | x_1 + x' \rangle$$

$$2\pi \delta(x') = \int \delta(x')$$

$$= \langle x_1 | \hat{\rho} | x_1 \rangle \checkmark$$

$$\int dx_1 W(x_1, x_2) = \int \frac{dx'}{\pi} e^{-2ix'x_2} \int dx_1 \langle x_1 - x' | \hat{\rho} | x_1 + x' \rangle$$

Aside $\int dx_1 \langle x_1 - x' | \hat{\rho} | x_1 + x' \rangle = \int dP dP' \int \langle x - x' | P \rangle \langle P' | x + x' \rangle \langle P | \hat{\rho} | P' \rangle dx_1$
insert complete moment set

$$= \int dP dP' \langle P | \hat{\rho} | P' \rangle \int \frac{dx_1}{2\pi} e^{i(x_1 - x')P} e^{-i(x_1 + x')P'}$$

(c) Suppose the operator is only a function of the general quadrature operators $\hat{X}(\theta) = \frac{\hat{a}e^{-i\theta} + \hat{a}^\dagger e^{i\theta}}{2}$ (i)

$$\begin{aligned}\hat{f} &\equiv f(\hat{X}_c(\theta)) = \sum_{n=0}^{\infty} \frac{1}{n!} \left. \frac{d^n f}{dx^n} \right|_0 (\hat{X}_c(\theta))^n \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} \left. \frac{d^n f}{dx^n} \right|_0 (\hat{a}e^{-i\theta} + \hat{a}^\dagger e^{i\theta})^n = \text{Symmetrically ordered product}\end{aligned}$$

$$\Rightarrow \langle \hat{f}_c \rangle = \int dX_1(\theta) dX_2(\theta) f(X_c(\theta)) W(X_1(\theta), X_2(\theta))$$

$\Rightarrow W(X_1(\theta), X_2(\theta))$ is the appropriate quasiprobability distribution when considering measurements of quadratures

Problem 3: The "Schrodinger cat" state

Given superposition of two coherent states:

$$|\psi\rangle = N(|\alpha_1\rangle + |\alpha_2\rangle) \quad N = \frac{1}{\sqrt{2(1 + e^{-|\alpha_1 - \alpha_2|^2})}}$$

Specialize to case $\alpha_1 = -\alpha_2 = \alpha_0$ (Real)

The Wigner function

$$W(\alpha) = \int \frac{d\beta}{\pi^2} \chi(\beta) e^{\alpha\beta^* - \alpha^*\beta}$$

$$\chi(\beta) = \text{Tr}(\hat{\rho} \hat{D}(\beta)) = \langle \psi | \hat{D}(\beta) | \psi \rangle$$

$$= N (\langle \alpha_0 | \hat{D}(\beta) | \alpha_0 \rangle + \langle -\alpha_0 | \hat{D}(\beta) | -\alpha_0 \rangle$$

$$+ \langle -\alpha_0 | \hat{D}(\beta) | \alpha_0 \rangle + \langle \alpha_0 | \hat{D}(\beta) | -\alpha_0 \rangle)$$

The first two terms represent the characteristic function of the coherent states $|\alpha_0\rangle$ and $|-\alpha_0\rangle$ respectively. $W(\alpha) = \frac{2}{\pi} e^{-2|\alpha \pm \alpha_0|^2}$

To calculate the off-diagonal terms, use part 1(c)

$$\langle -\alpha_0 | \hat{D}(\beta) | \alpha_0 \rangle = e^{-|\beta + 2\alpha_0|^2/2}$$

$$\int \frac{d\beta}{\pi^2} e^{-|\beta + 2\alpha_0|^2/2} e^{\alpha\beta^* - \alpha^*\beta}$$

$$\Rightarrow \text{let } \gamma = \beta + 2\alpha_0 \Rightarrow \beta = \gamma - 2\alpha_0$$

$$\int \frac{d\gamma}{\pi^2} e^{-\frac{|\gamma|^2}{2}} e^{\alpha\gamma^* - \alpha^*\gamma} e^{-2\alpha_0(\gamma - \alpha^*)} = \frac{2e^{-2|\alpha|^2}}{\pi} e^{-4i\alpha_0 \text{Im}(\alpha)}$$

$$= \frac{2}{\pi} e^{-2|\alpha|^2} \quad (\text{part 1a})$$

Putting it all together,

$$W(x, \lambda) = \frac{2N}{\pi} \left\{ e^{-2|x-x_0|^2} + e^{-2|x+x_0|^2} + 2e^{-2x^2} \cos(4x_0 \operatorname{Im}(\lambda)) \right\}$$

The Wigner function is negative for some x

\Rightarrow Nonclassical state

Let $\alpha = +iP$

$$\Rightarrow W(x, P) = \frac{2N}{\pi} \left\{ e^{-2\{(x-x_0)^2 + P^2\}} + e^{-2\{(x+x_0)^2 + P^2\}} + 2e^{-2(x^2 + P^2)} \cos(4x_0 P) \right\}$$

$$\text{Now } \int_{-\infty}^{\infty} dP W(x, P) = N \sqrt{\frac{2}{\pi}} \left(e^{-2(x-x_0)^2} + e^{-2(x+x_0)^2} \right) \\ \stackrel{?}{=} |\langle x | \psi \rangle|^2 = |\psi(x)|^2 + e^{-2(x^2 + x_0^2)}$$

Yes given $\langle x | \alpha_0 \rangle = \left(\frac{2}{\pi}\right)^{1/4} e^{-x^2}$ in our units

$$\int_{-\infty}^{\infty} dx W(x, P) = N \sqrt{\frac{2}{\pi}} \left(2e^{-2P^2} \right) \cos(4x_0 P)$$

$$= |\tilde{\psi}(P)|^2$$

\uparrow oscillation from superposition

Yes ✓

Following are plots of the Wigner function for $\alpha_0 = 1, 5, 10$. We see two kinds of feature

- (1) Gaussian packets localized near $\pm\alpha_0$
- (2) Oscillations as a function of P near $x=0$

Feature (1) is expected classically in terms of a distribution in phase space

Feature (2) ~~is~~ is the nonclassical interference coming from the "quantum superposition"

Note that as $\alpha_0 \rightarrow \infty$ (very nonclassically) the oscillations get more and more rapid.


This picture shows a kind of "interference in phase space". It captures both the position and momentum representations.

```
In[18]:= A = 2 e-2 x02 (1 + e2 x02)
```

```
Out[18]= 2 e-2 x02 (1 + e2 x02)
```


```
In[14]:= W[x_, p_] := 2 / (A Pi) (Exp[-2 ((x - x0)^2 + p^2)] +  
Exp[-2 ((x + x0)^2 + p^2)] + 2 Exp[-2 (x^2 + p^2)] Cos[4 x0 p])
```

```
In[84]:= x0 = 2;  
Plot3D[W[x, p], {x, -3, 3}, {p, -1.5, 1.5}, PlotRange -> All, PlotPoints -> 40,  
ViewPoint -> {1.409, -2.602, 0.320}]
```




```
Out[85]= - SurfaceGraphics -
```

```
In[88]:= x0 = 5;  
Plot3D[W[x, p], {x, -6, 6}, {p, -1.5, 1.5}, PlotRange -> All, PlotPoints -> 40,  
ViewPoint -> {1.409, -2.602, 0.320}]
```



```
Out[89]= - SurfaceGraphics -
```

```
In[100]:= x0 = 10;  
Plot3D[W[x, p], {x, -11, 11}, {p, -1.5, 1.5}, PlotRange -> All, PlotPoints -> 70,  
ViewPoint -> {1.409, -2.602, 0.320}]
```



```
Out[101]= - SurfaceGraphics -
```

Problem 4: Thermal Light

$$\hat{\rho} = \frac{1}{Z} e^{-\beta \hat{H}}$$

$$Z = \text{Tr}(e^{-\beta \hat{H}})$$

$$\hat{H} = \hbar \omega \hat{a}^\dagger \hat{a}$$

$$\Rightarrow Z = \sum_{n=0}^{\infty} (e^{-\beta \hbar \omega})^n = \frac{1}{1 - e^{-\beta \hbar \omega}}$$

$$\begin{aligned} (a) \langle n \rangle &= \frac{1}{Z} \sum_{n=0}^{\infty} n (e^{-\beta \hbar \omega})^n = \frac{1}{Z} \frac{d}{d(-\beta \hbar \omega)} \sum_{n=0}^{\infty} (e^{-\beta \hbar \omega})^n \\ &= \frac{1}{Z} \left(\frac{-e^{-\beta \hbar \omega}}{(1 - e^{-\beta \hbar \omega})^2} \right) = \frac{e^{-\beta \hbar \omega}}{1 - e^{-\beta \hbar \omega}} = \frac{1}{e^{\beta \hbar \omega} - 1} \end{aligned}$$

$$P_n = \frac{e^{-n \beta \hbar \omega}}{(1 - e^{-\beta \hbar \omega})^{-1}}$$

$$\begin{aligned} \text{now } e^{\beta \hbar \omega} &= 1 + \frac{1}{\langle n \rangle} \\ &= \frac{\langle n \rangle + 1}{\langle n \rangle} \end{aligned}$$

$$\Rightarrow P_n = \frac{\langle n \rangle^n}{(\langle n \rangle + 1)^n} \left[\frac{1}{1 - \frac{\langle n \rangle}{\langle n \rangle + 1}} \right]^{-1} = \frac{\langle n \rangle^n}{(1 + \langle n \rangle)^{n+1}}$$

$$\Delta n^2 = \langle n^2 \rangle + \langle n \rangle^2$$

$$\langle n^2 \rangle = - \frac{d}{d(\beta \hbar \omega)} \langle n \rangle \quad (\text{see e.g. Reif})$$

$$= \frac{+ e^{\beta \hbar \omega}}{(e^{\beta \hbar \omega} - 1)^2} = \frac{e^{\beta \hbar \omega} - 1 + 1}{(e^{\beta \hbar \omega} - 1)^2}$$

$$= \frac{1}{e^{\beta \hbar \omega} - 1} + \frac{1}{(e^{\beta \hbar \omega} - 1)^2}$$

$$= \langle n \rangle + \langle n^2 \rangle$$

↙ fluctuations beyond coherent state

$$\langle \hat{a} \rangle = \text{Tr}(\hat{\rho} \hat{a}) = \sum_n \langle n | \hat{a} | n \rangle \rho_{nn} = 0$$

\Rightarrow No mean amplitude in contrast to coherent state $|\alpha\rangle$.

(b) Calculating the quasi-distribution

The simplest is the Q-function (Husimi distribution)

$$\begin{aligned} Q(\alpha) &= \frac{1}{\pi} \langle \alpha | \hat{\rho} | \alpha \rangle = \frac{1}{2\pi} \langle \alpha | e^{-\beta \hat{H}} | \alpha \rangle \\ &= \frac{1}{2\pi} \langle \alpha | e^{-\beta \hbar \omega \hat{a}^\dagger \hat{a}} | \alpha \rangle \end{aligned}$$

Aside: Here's a good trick to remember

$$\langle \alpha | e^{-i\theta \hat{a}^\dagger \hat{a}} | \alpha \rangle = \langle \alpha | \alpha e^{-i\theta} \rangle$$

$$= \exp\left\{-(\alpha - \alpha e^{-i\theta})^2 / 2\right\} \exp\left\{\frac{|\alpha|^2 e^{-i\theta} - |\alpha|^2 e^{+i\theta}}{2}\right\}$$

$$= \exp\left\{-|\alpha|^2 (1 - e^{-i\theta})\right\}$$

This analytically continues $\theta \Rightarrow -i\beta\hbar\omega$

$$\Rightarrow \langle \alpha | e^{-\beta\hbar\omega \hat{a}^\dagger \hat{a}} | \alpha \rangle = \exp\left\{-|\alpha|^2 (1 - e^{-\beta\hbar\omega})\right\}$$

$= \frac{1}{2} - 1$

$$\therefore Q(\alpha) = \frac{1}{\pi^2} e^{-|\alpha|^2 / 2}$$

Given the Q-function, we can find P and W using the relationship between their characteristic functions.

$$\chi_Q(\beta) = \int dx Q(x) e^{\beta x^* - \beta^* x} = e^{-z/|\beta|^2} \quad (\text{using Problem 1a})$$

$$\chi_Q(\beta) = \text{Tr}(\hat{\rho} \hat{D}(\beta)) e^{-|\beta|^2/2}, \quad \chi_W(\beta) = \text{Tr}(\hat{\rho} \hat{D}(\beta))$$

$$\chi_P(\beta) = \text{Tr}(\hat{\rho} \hat{D}(\beta)) e^{+|\beta|^2/2}$$

$$\Rightarrow \chi_W(\beta) = e^{-(z-1/2)|\beta|^2} \quad \chi_P(\beta) = e^{-(z-1)|\beta|^2}$$

Aside $z-1 = \frac{1}{1-e^{-\beta \hbar \omega}} - 1 = \frac{1}{e^{\beta \hbar \omega} - 1} = \langle n \rangle$

$$\Rightarrow \chi_W(\beta) = e^{-(\langle n \rangle + 1/2)|\beta|^2} \quad \chi_P(\beta) = e^{-\langle n \rangle |\beta|^2}$$

$$W(x) = \frac{1}{\pi(\langle n \rangle + 1/2)} e^{-|x|^2/(\langle n \rangle + 1/2)}$$

$$P(x) = \frac{1}{\pi \langle n \rangle} e^{-|x|^2/\langle n \rangle}$$

$$Q(x) = \frac{1}{\pi(\langle n \rangle + 1)} e^{-|x|^2/(\langle n \rangle + 1)}$$

All Gaussian functions, with slightly different widths.

In the limit $\langle n \rangle \rightarrow 0$

$$W(\alpha) \rightarrow \frac{2}{\pi} e^{-2|\alpha|^2} = \text{Wigner function for a vacuum}$$

$$P(\alpha) \rightarrow \delta^{(2)}(\alpha) = \text{P function for vacuum}$$

$$Q(\alpha) \rightarrow \frac{1}{\pi} e^{-|\alpha|^2} = \frac{1}{\pi} |\langle 0|\alpha\rangle|^2 = \text{Q function for vacuum}$$

In the view of quasiprobability the thermal state and vacuum are very similar. They are both Gaussian states centered at the origin. The width of the Gaussian depends on the "temperature"; i.e. $\langle n \rangle$, and the particular operator ordering used. The Q and W functions include quantum fluctuations; their widths never go to zero.

The P function describes the state as a statistical mixture of coherent states. At zero temperature ($\langle n \rangle = 0$) this distribution collapses to a delta function.

(c) To calculate $\Delta X_1^2(\theta)$ and $\Delta X_2^2(\theta)$ we use the Wigner function (Problem 2). These variances are just the variance of the Wigner Gaussian (independent of θ)

$$\Delta X_1^2 = \Delta X_2^2 = \langle n \rangle + \frac{1}{2} \rightarrow \frac{1}{2} \text{ for } \langle n \rangle = 0$$

To calculate $\Delta n^2 = \langle n^2 \rangle - \langle n \rangle^2$

$$\text{Recall } \langle n^2 \rangle = \langle a^\dagger a a^\dagger a \rangle = \langle a^{\dagger 2} a^2 \rangle + \langle n \rangle$$

$$\Rightarrow \Delta n^2 = \langle :n^2: \rangle - \langle :n: \rangle^2 + \langle n \rangle$$

$$= (\Delta |x|^2)^2 + \langle n \rangle \leftarrow \text{"shot noise"}$$

\uparrow
Uncertainty in coherent state intensity from P-function

$$\Rightarrow \Delta n^2 = \langle n^2 \rangle + \langle n \rangle$$

Photon bunching!

The fluctuations in Δn^2 have two components

- "shot noise" arising from the particle nature
- Statistical fluctuation in $|x|^2$ for a finite temperature.