

Quantum information theory

Lectures 9-10

Quantum states. I-II. Mixed states.

Pure states:

$$\text{State vector } |\psi\rangle = \sum_j |e_j\rangle \underbrace{\langle e_j | \psi \rangle}_{c_j = (\text{probability amplitude})}$$

$$1 = \langle \psi | \psi \rangle = \sum_j |c_j|^2$$

Measurement in basis $|e_j\rangle$:

$$P_k = |\langle e_k | \psi \rangle|^2 = \langle e_k | \psi \rangle \langle \psi | e_k \rangle = \langle e_k | P_\psi | e_k \rangle$$

Mixed states: What if we know system is in state $|\psi_j\rangle$ with probability g_j (ensemble)

$$P_k = \sum_j \underbrace{P_{kj}}_{|\langle e_k | \psi_j \rangle|^2} g_j = \langle e_k | \left(\sum_j g_j |\psi_j\rangle \langle \psi_j| \right) | e_k \rangle$$

$$|\langle e_k | \psi_j \rangle|^2 = \langle e_k | \psi_j \rangle \langle \psi_j | e_k \rangle$$

$$P_k = \text{tr}(\rho |e_k\rangle \langle e_k|)$$

$$\text{Observable } A = \sum_k \lambda_k |e_k\rangle \langle e_k|$$

$$\langle A \rangle = \sum_k \lambda_k P_k = \text{tr}(\rho A)$$

density operator ρ

$\{g_j, |\psi_j\rangle\}$ or $\{\sqrt{g_j}, |\psi_j\rangle\}$ is an ensemble decomposition

of ρ .

[A density operator is a Hermitian operator with nonnegative eigenvalues that sum to 1. ($\text{tr} \rho = 1$).]

Let $|\phi_j\rangle$, $j=1, \dots, D$ be a basis (i.e., D linearly independent, not necessarily orthogonal or normalized) for a D -dimensional Hilbert space. Any vector can be expanded uniquely as

$$|\psi\rangle = \sum_j c_j |\phi_j\rangle$$

If $|e_n\rangle$, $n=1, \dots, D$, is an orthonormal basis,

$$\langle e_n | \psi \rangle = \sum_j c_j \underbrace{\langle e_n | \phi_j \rangle}_{S_{nj}}$$

$$\begin{pmatrix} \langle e_1 | \psi \rangle \\ \vdots \\ \langle e_D | \psi \rangle \end{pmatrix} = S \begin{pmatrix} c_1 \\ \vdots \\ c_D \end{pmatrix}$$

S is invertible because of the uniqueness of the c_j (i.e., the l.i. of $|\phi_j\rangle$).

$$\langle \phi_j | \psi \rangle = \sum_k \underbrace{\langle \phi_j | e_n \rangle}_{\langle e_n | \phi_j \rangle^* = S_{nj}^* = (S^{\dagger})_{jk}} \langle e_n | \psi \rangle \quad (S^{\dagger})^{-1} = (S^{-1})^{\dagger}$$

$$\begin{pmatrix} \langle \phi_1 | \psi \rangle \\ \vdots \\ \langle \phi_D | \psi \rangle \end{pmatrix} = S^{\dagger} \begin{pmatrix} \langle e_1 | \psi \rangle \\ \vdots \\ \langle e_D | \psi \rangle \end{pmatrix} = S^{\dagger} S \begin{pmatrix} c_1 \\ \vdots \\ c_D \end{pmatrix}$$

$$\begin{pmatrix} \langle e_1 | \psi \rangle \\ \vdots \\ \langle e_D | \psi \rangle \end{pmatrix} = (S^{-1})^{\dagger} \begin{pmatrix} \langle \phi_1 | \psi \rangle \\ \vdots \\ \langle \phi_D | \psi \rangle \end{pmatrix}$$

These inner products specify $|\psi\rangle$.

Back up 2 steps:

Hermitian operators are a real vector space; L_V is the complexification. \textcircled{P}

1. The operators on a D -dimensional Hilbert space V make up a D^2 -dimensional complex vector space L_V , with inner product $(A, B) = \text{tr}(A^\dagger B)$.

Orthonormal basis for L_V : $\tau_{jk} = |e_j\rangle\langle e_k|$



$$A = \sum_{j,k} A_{jk} |e_j\rangle\langle e_k|$$

$$\begin{aligned} \text{tr}(\tau_{em}^\dagger \tau_{jk}) &= \text{tr}(|e_m\rangle\langle e_l| |e_j\rangle\langle e_k|) \\ &= \langle e_l | e_m \rangle \langle e_l | e_j \rangle \\ &= \delta_{jl} \delta_{km} \end{aligned}$$

There is no such basis in a real vector space.

Basis of 1-d projectors: (pure states)

This is not an orthonormal basis. That's impossible

$$\text{tr}(|\psi\rangle\langle\psi| |\psi\rangle\langle\psi|) = \langle\psi|\psi\rangle^2$$

$$\left. \begin{array}{l} |\phi_a\rangle \\ a=1, \dots, D^2 \end{array} \right\} \begin{array}{l} j=1, \dots, D: |e_j\rangle\langle e_j| = \tau_{jj} \\ j < k: |X_{jk}\rangle\langle X_{jk}| = \frac{1}{2}(\tau_{jj} + \tau_{kk} + \tau_{jk} + \tau_{kj}) \\ |E_{jk}\rangle\langle E_{jk}| = \frac{1}{2}(\tau_{jj} + \tau_{kk} + i(-\tau_{jk} + \tau_{kj})) \end{array}$$

$$|X_{jk}\rangle = \frac{1}{\sqrt{2}}(|e_j\rangle + |e_k\rangle)$$

$$|E_{jk}\rangle = \frac{1}{\sqrt{2}}(|e_j\rangle + i|e_k\rangle)$$

An operator A is specified by the inner products

$$(|\phi_a\rangle\langle\phi_a|, A) = \text{tr}(|\phi_a\rangle\langle\phi_a| A) = \langle\phi_a| A |\phi_a\rangle$$

An operator A is (over) specified by

$\langle \psi | A | \psi \rangle$ for all $|\psi\rangle$.

"sandwiches"

An operator A is Hermitian iff $\langle \psi | A | \psi \rangle$ is real for all $|\psi\rangle$.

② A positive operator G is one such that $\langle \psi | G | \psi \rangle$ is real and nonnegative for all $|\psi\rangle$.

This is denoted $G \geq 0$. An operator G

is positive iff it is Hermitian with nonnegative eigenvalues.

Positive operators G_1 and G_2 are orthogonal, i.e. $\text{tr}(G_1 G_2) = 0$, iff $G_1 G_2 = 0$.

Positive definite $G > 0$ means $\langle \psi | G | \psi \rangle > 0$ for all $|\psi\rangle$.

\iff (i) Hermitian with positive eigenvalues

(ii) Positive and invertible.

Positive operators have a square root:
 $\sqrt{G} = \sum_i \sqrt{\lambda_i} |e_i\rangle\langle e_i|$.

Characterization of a density operator. ρ is a density operator iff (i) $\rho \geq 0$ and (ii) $\text{tr}(\rho) = 1$.

Pure states: $\rho = |\psi\rangle\langle\psi|$ is a 1-d (rank-1) projector.

① A unit-trace Hermitian operator ρ is a pure state

iff $\rho^2 = \rho$. [A Hermitian operator P is a projector iff $P^2 = P$.
 (11) squares to itself, but is not a projector.]

② A density operator ρ is a pure state iff $\text{tr}(\rho^2) = 1$.

\iff : trivial

\implies : $0 = \text{tr}(G_1 G_2) = \text{tr}(G_1^{1/2} G_1^{1/2} G_2^{1/2} G_2^{1/2}) = \text{tr}((G_1^{1/2} G_2^{1/2})^\dagger G_1^{1/2} G_2^{1/2})$

$\implies 0 = G_1^{1/2} G_2^{1/2} \implies 0 = G_1 G_2$

Convex sets: A set S is convex if for any $v_1, v_2 \in S$,


$$\lambda v_1 + (1-\lambda)v_2 \in S, \quad 0 \leq \lambda \leq 1.$$



Examples:

$$\sum_i \lambda_i v_i \in S, \quad \lambda_i \geq 0, \quad \sum_i \lambda_i = 1$$

convex combination or mixture



An extreme point of S is a point that can't be written as a proper convex combination (0 < λ < 1) of any other points.

If S is closed and bounded, every point in S can be written as a convex combination of extreme points.

Simplex: Closed & bounded convex set where expansion in terms of extreme points is unique.

[The density operators are a closed and bounded convex set whose extreme points are the pure states.]

Qubits:

$$\rho = A_0 \mathbb{1} + \vec{A} \cdot \vec{\sigma} = (A_0 + |\vec{A}|) |\vec{n}\rangle\langle\vec{n}| + (A_0 - |\vec{A}|) |\vec{-n}\rangle\langle\vec{-n}|$$


Hermitian $\Rightarrow A_0, \vec{A}$ real

positive $\Rightarrow |\vec{A}| \leq A_0 = 1/2$

$\mathbb{1} = \text{tr}(\rho) \Rightarrow A_0 = 1/2$

$$\rho = \frac{1}{2} (\mathbb{1} + \vec{s} \cdot \vec{\sigma}), \quad |\vec{s}| \leq 1$$
$$\vec{s} = \text{tr}(\rho \vec{\sigma}) = \begin{pmatrix} \text{Bloch} \\ \text{vector} \end{pmatrix}$$

Bloch sphere



Pure states on surface

Mixed states in interior

Contrast higher dimensions:

Mixed states: (d^2-1) -dimensional manifold

Pure states: $2(d-1)$ dimensional

Multiple boundaries for every number of zero eigenvalues from 1, ..., $d-1$.

Freedom in ensemble decompositions for qubits:

$$\rho = \sum_j p_j |\psi_j\rangle\langle\psi_j| = \frac{1}{2} \left(1 + \vec{\sigma} \cdot \underbrace{\sum_j p_j \vec{\sigma}_j}_{\vec{S}} \right)$$

$$\frac{1}{2} (1 + \vec{\sigma} \cdot \vec{S})$$

Freedom in ensemble decompositions (HJW theorem;

Schrödinger). Space of density operators is not a simplex.

The rank of ρ is the dimension of the support

Eigendecomposition $\rho = \sum_{j=1}^D \lambda_j |e_j\rangle\langle e_j| = \sum_{j=1}^D |\bar{e}_j\rangle\langle \bar{e}_j|$

$$|\bar{e}_j\rangle = \sqrt{\lambda_j} |e_j\rangle$$

Support S of ρ is the subspace spanned by eigenvectors with nonzero eigenvalue. Null subspace

(kernel) K of ρ is the subspace spanned by eigenvectors with zero eigenvalue. These subspaces are orthogonal. The subspace orthogonal to a subspace S is called the orthocomplement of S .

$$P_S = \sum_{j \neq 0} |e_j\rangle\langle e_j|$$

$$P_R = \sum_{j=0} |e_j\rangle\langle e_j|$$

Ensemble decomposition: $\{P_\alpha, |\psi_\alpha\rangle\}$ or $\{|\bar{\psi}_\alpha\rangle = \sqrt{P_\alpha}|\psi_\alpha\rangle\}$

$$\rho = \sum_\alpha P_\alpha |\psi_\alpha\rangle\langle\psi_\alpha| = \sum_\alpha |\bar{\psi}_\alpha\rangle\langle\bar{\psi}_\alpha|$$

All states in the ensemble decomposition lie in the support of ρ [Proof: If $|e_j\rangle$ lies in the null subspace, then $0 = \langle e_j|\rho|e_j\rangle = \sum_\alpha |\langle e_j|\bar{\psi}_\alpha\rangle|^2 \Rightarrow \langle e_j|\bar{\psi}_\alpha\rangle = 0$ for all $\alpha \Rightarrow |\bar{\psi}_\alpha\rangle$ is orthogonal to the null subspace and thus lies in the support.]

HJW theorem. Two ensemble decompositions, $\{|\bar{\psi}_\alpha\rangle\}$ and $\{|\bar{\phi}_\alpha\rangle\}$, correspond to the same density operator iff there exists a unitary matrix

$U_{\alpha\beta}$ such that

$$|\bar{\phi}_\alpha\rangle = \sum_\beta U_{\alpha\beta} |\bar{\psi}_\beta\rangle$$

Add zero vectors to ensemble with fewer elements.

Column-vector notation

Proof:

$$\begin{aligned} \sum_\alpha |\bar{\phi}_\alpha\rangle\langle\bar{\phi}_\alpha| &= \sum_{\alpha, \beta, \gamma} U_{\alpha\beta} |\bar{\psi}_\beta\rangle\langle\bar{\psi}_\gamma| U_{\alpha\gamma}^* \\ &= \sum_{\beta, \gamma} |\bar{\psi}_\beta\rangle\langle\bar{\psi}_\gamma| \underbrace{\sum_\alpha U_{\alpha\beta} U_{\alpha\gamma}^*}_{\delta_{\beta\gamma}} \\ &= \sum_\beta |\bar{\psi}_\beta\rangle\langle\bar{\psi}_\beta| \end{aligned}$$

→
$$P = \sum_{\alpha} |\bar{\psi}_{\alpha}\rangle \langle \bar{\psi}_{\alpha}| = \sum_{|e_j\rangle \in S} |e_j\rangle \langle e_j|$$

↑
eigendecomposition

We know $|e_j\rangle \in S$, so this sum can be restricted to the support

$$|\bar{\psi}_{\alpha}\rangle = \sum_{|e_j\rangle \in S} |e_j\rangle \langle e_j | \bar{\psi}_{\alpha}\rangle = \sum_{|e_j\rangle \in S} |e_j\rangle \underbrace{\langle e_j | \bar{\psi}_{\alpha}\rangle}_{M_{\alpha j}}$$

← $\lambda_j \neq 0$ because $|e_j\rangle \in S$

$$\begin{aligned} \sum_{\alpha} M_{\alpha j} M_{\alpha k}^* &= \left\langle e_j \left| \frac{\sum_{\alpha} |\bar{\psi}_{\alpha}\rangle \langle \bar{\psi}_{\alpha}|}{\sqrt{\lambda_j \lambda_k}} \right| e_k \right\rangle \\ &= \frac{\langle e_j | P | e_k \rangle}{\sqrt{\lambda_j \lambda_k}} \\ &= \frac{\lambda_j \delta_{jk}}{\sqrt{\lambda_j \lambda_k}} \\ &= \delta_{jk} \end{aligned}$$

$M_{\alpha j}$ is a $N \times R$ matrix, where N is the number of elements in the ensemble decomposition and R is the rank of P .

M can be extended to be unitary.