

**6.1 Complete positivity of unital trace-preserving qubit quantum operations.** An arbitrary trace-preserving qubit quantum operation can be written as

$$\mathcal{A}(\rho) = V\mathcal{B}(U\rho U^\dagger)V^\dagger,$$

where

$$\mathcal{B}(\rho) = \frac{1}{2}(1 + \boldsymbol{\sigma} \cdot (T\mathbf{S} + \mathbf{d})),$$

and where

$$T = \begin{pmatrix} t_x & 0 & 0 \\ 0 & t_y & 0 \\ 0 & 0 & t_z \end{pmatrix}$$

is a diagonal matrix that contracts (and perhaps inverts or reflects, depending on the number of negative diagonal entries) the Bloch sphere and  $\mathbf{d}$  is a displacement of the contracted sphere. The initial and final unitaries,  $U$  and  $V$ , represent initial and final rotations of the Bloch sphere. Since these unitary transformations are completely positive, the complete positivity of  $\mathcal{A}$  is determined by the complete positivity of  $\mathcal{B}$ .

In this problem we deal with the special case where there is no displacement of the Bloch sphere, i.e.,  $\mathbf{d} = 0$ . In this case the operation takes the maximally mixed state to itself, i.e.,  $\mathcal{B}(1) = 1$  (such a quantum operation is called *unital*). Positivity of  $\mathcal{B}$  requires the output states to be in the Bloch sphere, which means that the diagonal entries of  $T$  must satisfy  $|t_j| \leq 1$ . Thus positivity alone allows the diagonal entries to lie anywhere within a cube of side length 2 centered at the origin. The goal of this problem is to explore the further restrictions imposed by complete positivity.

(a) Find the restrictions imposed by complete positivity, and describe or draw the region of allowed values of  $t_x$ ,  $t_y$ , and  $t_z$ .

(b) What kinds of maps of the Bloch sphere are allowed by complete positivity when one axis is left unchanged, say,  $t_z = 1$ ?

(c) The unital qubit quantum operations form a convex set. What are the extreme points of this set?