

## Physics 511 Spring 2000

### Problem Set #2: DUE Wed. Feb. 1


**Read: Jackson Chap. 4, Low Chap. 1**  
**Supplementary notes on Multipoles**

#### (1) The "Yukawa Potential" (10 Points)

The electric force dominates the interaction of particles because it is very long-range; i.e. the electrostatic potential falls off as an algebraic function of distance  $1/r$ . In certain circumstances we would like to "cut-off" the Coulomb field after some distance. This can be achieved using the "Yukawa" potential.

$$\Phi(r) = G \frac{ae^{-r/a}}{r},$$

where  $G$  is some coupling constant, and  $a$  is a distance. This potential can be mimicked in electrostatics by a point charge  $q$  at the origin and "screened" by a smeared out negative charge distribution:  $\rho(\mathbf{x}) = -q\delta^{(3)}(\mathbf{x}) + \frac{q}{4\pi a^2} \frac{e^{-r/a}}{r}$ . (Here  $a$  is a constant with dimensions of length)




- Sketch  $\Phi(r)$  showing the asymptotic form for  $r \ll a$  and  $r \gg a$ . Explain the physical significance of the constant  $a$ .
- Show that the overall charge distribution is neutral. Does this make sense given the long range form of the potential?
- Derive the Yukawa potential for this charge distribution.

(Hint Use Gauss' law to find the electric field first)

- What is the potential energy stored in this charge distribution? Explain your result.

**(2) Potentials and contours (15 points)**

An infinitely long strip of width  $L$  (shown below) carries a charge per unit area  $\sigma$



(a) Find the electrostatic potential and the electric field in the  $x$ - $y$  plane.

(b) Take the following limits of  $\mathbf{E}$  and  $\Phi$ , and explain why the result is what you expect.

(i)  $\lim \frac{R}{L} \rightarrow 0$  ,                      (ii)  $\lim \frac{R}{L} \rightarrow \infty$  ,                      where  $R = \sqrt{x^2 + y^2}$

(c) Plot the field and equipotential contours. Do this 3 times with different ranges for the plots: (i)  $x, y \sim L$ . (ii)  $x, y \gg L$ . (iii)  $x, y \ll L$ . Explain your Results


(d) Now suppose there are two strips, one with positive surface charge  $\sigma$  at  $y=0$ , and one with equal and opposite charge  $-\sigma$ , at  $y = -s$  Use the *principle of superposition* to find  $\Phi$ ,  $\mathbf{E}$

(e) Again plot the contours and electric field. Do this for two cases

(i)  $s \ll L$ . (ii)  $s \gg L$  Explain your results.

**(3) Boundary condition at a surface charge (10 points)**

(a) Each of the following geometries has a uniform charge density  $\rho$ . Find the electric field every in space. Sketch the plot of  $E$  as a function of the appropriate distance.



- (i) Spherical shell, inner radius  $a$ , outer radius  $b$ ,  
 (ii) Cylindrical shell, length  $L \gg a, b$   
 (iii) Infinite slab, thickness  $d$

(b) Take the limit where thickness of each shell(slab) goes to zero, such that all of the charge is concentrated on a surface with density  $\sigma$ ; i. e.,

- (i)  $b - a \rightarrow 0$ ,  $\rho(b - a) \rightarrow \sigma$   
 (ii)  $b - a \rightarrow 0$ ,  $\rho(b - a) \rightarrow \sigma$   
 (iii)  $d \rightarrow 0$ ,  $\rho d \rightarrow \sigma$

Show that the electric field normal to the surface is discontinuous by  $\Delta E_{\perp} = 4\pi\sigma$

**(4) The Dirac Delta Function: Jackson 1.3 (10 points)**