

## Physics 531

### Problem Set #4

Due Thursday, Oct. 13, 2011

#### Problem 1: Landé Projection Theorem

The Landé Projection Theorem (LPT) is a special case of the Wigner-Eckart Theorem for the case that the initial and final state are the same angular momentum. It states:

$$\langle \alpha'; j m' | \hat{V} | \alpha; j m \rangle = \frac{\langle \alpha'; j | \hat{\mathbf{J}} \cdot \hat{V} | \alpha; j \rangle}{j(j+1)} \langle j m' | \hat{\mathbf{J}} | j m \rangle, \text{ where } \hat{V} \text{ is a vector operator w.r.t. } \hat{\mathbf{J}}.$$

- (a) Give a geometric interpretation of this in terms of a vector picture.  
(b) To prove this theorem, take the following steps (do not give verbatim, Sakurai's derivation):
- Show that  $\langle \alpha'; j, m | \hat{\mathbf{J}} \cdot \hat{V} | \alpha; j, m \rangle = \langle \alpha'; j | J | \alpha'; j \rangle \langle \alpha'; j | V | \alpha; j \rangle$ , independent of  $m$ .
  - Use this to show,  $\langle \alpha; j | J | \alpha; j \rangle^2 = j(j+1)$  independent of  $\alpha$ .
  - Show that  $\langle j m' | J_q | j m \rangle = \langle j m' | \hat{J}_q | j m \rangle / \sqrt{j(j+1)}$ .
  - Put it all together to prove the LPT.

- (c) As an application of the LPT, consider the Zeeman interaction for hydrogen.

$$\hat{H}_{\text{int}} = -\hat{\boldsymbol{\mu}} \cdot \mathbf{B},$$

where the magnetic dipole operators is  $\hat{\boldsymbol{\mu}} = -\mu_B(\hat{\mathbf{L}} + 2\hat{\mathbf{S}})$ , ignoring the nuclear magnetic moment.

If the Zeeman effect is small compared to the fine-structure, but ignoring hyperfine structure, use the LPT to show that in a state  $nL_j$ , the magnetic moment has the form,

$$\hat{\boldsymbol{\mu}} = -g_j \mu_B \hat{\mathbf{J}}, \text{ where } g_j = \frac{3}{2} + \frac{S(S+1) - L(L+1)}{2J(J+1)} \text{ is known as the Landé g-factor.}$$

$$\text{Hint: Use } \mathbf{J} \cdot \mathbf{L} = \mathbf{L}^2 + \frac{1}{2}(\mathbf{J}^2 - \mathbf{L}^2 - \mathbf{S}^2) \text{ and } \mathbf{J} \cdot \mathbf{S} = \mathbf{S}^2 + \frac{1}{2}(\mathbf{J}^2 - \mathbf{L}^2 - \mathbf{S}^2).$$

- (d) Estimate the magnetic field strength at which the Zeeman interaction becomes on the order of the fine structure splitting between the  $2p_{1/2}$  and  $2p_{3/2}$  state in hydrogen.

- (e) Now solve for the energy shift of the sublevels exactly, including both the spin-orbit and Zeeman interaction, as in P.S.02, #3. Show that for small  $\mathbf{B}$  fields, one arrives at the "linear Zeeman shift" found in part (c).

**Problem 2: Natural lifetimes of Hydrogen** (10 points)

Though in the absence of any perturbation, an atom in the excited state will stay there forever (it is a stationary state), in reality it will “spontaneously decay” to the ground state. Fundamentally this occurs because the atom is always perturbed by ‘vacuum fluctuations’ in the electro-magnetic field. The spontaneous emission rate on a dipole allowed transition from initial excited state  $|\psi_e\rangle$  to all allowed ground states  $|\psi_g\rangle$  is ,

$$\Gamma = \frac{4}{3\hbar} k^3 \sum_g \left| \langle \psi_g | \hat{\mathbf{d}} | \psi_e \rangle \right|^2, \text{ where } k = \omega_{eg} / c \text{ is the emitted photon's wave number.}$$

Consider now hydrogen including fine-structure. For a given sublevel, the spontaneous emission rate is

$$\Gamma_{(nLJM_J) \rightarrow (n'L'J'M'_J)} = \frac{4}{3\hbar} k^3 \sum_{M'_J} \left| \langle n'L'J'M'_J | \hat{\mathbf{d}} | nLJM_J \rangle \right|^2.$$

(a) Show that the spontaneous emission rate is *independent* of the initial  $M_J$ . Explain this result physically.

(b) Calculate the lifetime ( $\tau=1/\Gamma$ ) of the  $2p_{1/2}$  state in seconds.

**Problem 3: Three spherical harmonics**

As we have seen many times, often we need to calculate an integral of the form,

$$\int d\Omega Y_{l_3 m_3}^*(\theta, \phi) Y_{l_2 m_2}(\theta, \phi) Y_{l_1 m_1}(\theta, \phi).$$

This can be interpreted as the matrix element  $\langle l_3 m_3 | \hat{Y}_{m_2}^{(l_2)} | l_1 m_1 \rangle$ , where  $\hat{Y}_{m_2}^{(l_2)}$  is an irreducible tensor operator.

(a) Use the Wigner-Eckart theorem to determine the restrictions on the quantum numbers so that this integral does not vanish.

(b) Given the “addition rule” for Legendre polynomials:

$$P_{l_1}(\mu) P_{l_2}(\mu) = \sum_{l_3} \langle l_3 0 | l_1 0 l_2 0 \rangle^2 P_{l_3}(\mu),$$

Use the Wigner-Eckart theorem to prove

$$\int d\Omega Y_{l_3 m_3}^*(\theta, \phi) Y_{l_2 m_2}(\theta, \phi) Y_{l_1 m_1}(\theta, \phi) = \sqrt{\frac{(2l_2+1)(2l_1+1)}{4\pi(2l_3+1)}} \langle l_3 0 | l_2 0 l_1 0 \rangle \langle l_3 m_3 | l_2 m_2 l_1 m_1 \rangle$$

Hint: Consider  $\langle l_3 0 | \hat{Y}_0^{(l_2)} | l_1 0 \rangle$