

Lecture 5b: Two-level atomic response

- "Natural Lineshape"

Recall the time evolution of mean dipole-moment ("Lab Frame")

$$\langle \vec{d}(t) \rangle = \vec{d}_{\text{eq}} (\bar{u}(t) \cos \omega t - \bar{v}(t) \sin \omega t)$$

$\bar{u}(t) \Rightarrow$ In phase with $E \Rightarrow$ Index of refraction

$-\bar{v}(t) \Rightarrow$ In quadrature with $E \Rightarrow$ Absorption/ emission

$$\frac{\bar{u} + i\bar{v}}{2} = \rho_{ge}$$

Steady state \Rightarrow After transients decay $\Rightarrow \dot{\rho} = 0$

\Rightarrow Steady state coherences $\dot{\rho}_{ge} = 0$

$$\Rightarrow 0 = \left(-i\Delta - \frac{\Gamma}{2} \right) \rho_{ge}^{\text{s.s.}} + i \frac{\Omega}{2} (\rho_{ee}^{\text{s.s.}} - \rho_{gg}^{\text{s.s.}})$$

$$\Rightarrow \boxed{\rho_{ge}^{\text{s.s.}} = \left(\frac{\Omega/2}{\Delta - i\frac{\Gamma}{2}} \right) (\rho_{ee}^{\text{s.s.}} - \rho_{gg}^{\text{s.s.}})}$$


Suppose in steady state $\rho_{ee}^{\text{s.s.}} \approx 0$ (We come back to see when true)
 $\Rightarrow \rho_{gg} \approx 1$

\Rightarrow For weak excitation

$$\boxed{\rho_{ge}^{\text{s.s.}} = \left(\frac{-\Omega/2}{\Delta - i\frac{\Gamma}{2}} \right)}$$

"Complex Lorentzian"

"Linear response" \Rightarrow Classical S.I.F.O.



$$\vec{d}(+) = \text{Re}(\tilde{\chi} \tilde{E}_0 e^{-i\omega t})$$



↑
polarizability

$$\langle \vec{d}(t) \rangle_{ss.} = \text{Re} \left[\vec{d}_{\text{deg}} \underbrace{(u^{ss.} - i v^{ss.}) e^{-i\omega t}}_{\star} \right]$$

$$\Rightarrow 2 \vec{d}_{\text{deg}}^{ss.} = \frac{-\Omega}{\Delta + i \frac{\Gamma}{2}} = -\frac{\vec{d}_{\text{deg}} \cdot \tilde{E}}{\hbar (\Delta + i \frac{\Gamma}{2})}$$

\therefore Quantum polarizability

$$\tilde{\chi} = \frac{-|\vec{d}_{\text{deg}}|^2}{\hbar (\Delta + i \frac{\Gamma}{2})} = \frac{+|\vec{d}_{\text{deg}}|^2}{\hbar} \left(\frac{-\Delta + i \frac{\Gamma}{2}}{\Delta^2 + \frac{\Gamma^2}{4}} \right)$$



\nearrow
Lorentzian

Natural Line width = Γ

\Rightarrow Absorption strong when $\Delta < \Gamma$

Rate equations

In the regime when the transient dynamics have damped out and the coherences reach steady state, we can look at population dynamics.

Plugging in the coherences ~~here~~ is S.S.

$$\Rightarrow \dot{\rho}_{ee} = -\Gamma_{Pee} - \Omega \operatorname{Im}(\rho_{ge}^{\text{s.s.}})$$

$$\dot{\rho}_{ee} = -\Gamma_{Pee} + \left(\frac{\Omega^2/4}{\Delta^2 + \frac{\Gamma^2}{4}} \right) (\rho_{ee} - \rho_{gg})$$

Interpretation : Population rate equations

$$N_e \equiv \rho_{ee} \quad N_g \equiv \rho_{gg}$$

$$\left(\frac{\Omega^2/4}{\Delta^2 + \frac{\Gamma^2}{4}} \right) \Gamma = \Gamma_{\text{stim}} = \begin{aligned} & \text{stimulated absorption} \\ & \text{= stimulated emission} \end{aligned}$$

Check: Fermi's Golden rule:

$$\Gamma_{\text{stim}} = \frac{2\pi}{\hbar^2} |\langle \text{el } \hat{H}_{\text{int}}^{(+)} | g \rangle|^2 D(\omega_L)$$

$D(\omega_L)$ density of states

$$\hat{H}_{\text{int}}^{(+)} = -\frac{\hbar\Omega}{2} \hat{J}_+ e^{-i\omega_L t}$$


$$D(\omega_L) = \text{atomic line shape (normalized)} = \frac{\Gamma/2\pi}{\Delta^2 + \frac{\Gamma^2}{4}}$$

$$\Rightarrow \Gamma_{\text{stim}} = \frac{2\pi}{\hbar^2} \left(\frac{\hbar\Omega^2}{4} \right) \frac{\Gamma/2\pi}{\Delta^2 + \frac{\Gamma^2}{4}} = \frac{\frac{\Omega^2\Gamma}{4}}{\Delta^2 + \frac{\Gamma^2}{4}} \checkmark$$

⇒ Rate equations

$$\dot{N}_e = -(\Gamma + \Gamma_{\text{stim}}) N_e + \Gamma_{\text{stim}} N_g$$

$$\dot{N}_g = -\Gamma_{\text{stim}} N_g + (\Gamma + \Gamma_{\text{stim}}) N_e$$



$$\dot{N}_e = -\dot{N}_g$$

$$N_e + N_g = 1$$

Steady state ⇒ detailed balance $\dot{N}_e = \dot{N}_g = 0$

⇒ Steady state populations

$$N_e = 0 = -(\Gamma + \Gamma_{\text{stim}}) N_e^{\text{s.s.}} + \Gamma_{\text{stim}} (1 - N_e^{\text{s.s.}})$$

$$\Rightarrow N_e^{\text{s.s.}} = \frac{\Gamma_{\text{stim}}}{\Gamma + 2\Gamma_{\text{stim}}} = \frac{S/2}{1+S}$$

$$N_g^{\text{s.s.}} = \frac{1+S/2}{1+S}$$

Here I have defined:

Saturation parameter:

$$S \equiv \frac{2\Gamma_{\text{stim}}}{\Gamma} = \frac{S^2/2}{\Delta^2 + \frac{\Gamma^2}{4}}$$

For a given detuning and "oscillator strength", the saturation parameter determines the intensity ($\sim S^2$) at which we pump substantial population into the excited states. When $S \ll 1$

$$N_g \approx 1 - S/2, \quad N_e \approx S/2 \Rightarrow \text{Linear regime}$$

when $s \rightarrow \infty$, $N_e^{s.s.} \approx N_g \approx \frac{1}{2}$

\Rightarrow the transition is "saturated".

Resonant behavior: $\Delta = 0$

$$S_0 = \frac{2\Omega^2}{\Gamma^2} \Rightarrow s = 1 \text{ when } \Omega = \frac{\Gamma}{\sqrt{2}}$$

Since $\Omega = \frac{deg E}{\hbar} \propto E \Rightarrow \Omega^2 \propto E^2 \propto I$ (Intensity)

$$\Rightarrow S_0 = \frac{I}{I_{sat}} \text{ where } I_{sat} = \text{saturation intensity}$$

$$I_{sat} = \frac{I}{S_0} = \frac{I}{\frac{2\Omega^2}{\Gamma^2}} = \frac{\hbar^2 \Gamma^2}{2 deg} \frac{I}{E^2} = \left(\frac{\hbar^2 \Gamma^2}{2 deg} \right) \left(\frac{C E^2}{8\pi} \right)$$

$$I_{sat} = \frac{C}{16\pi} \frac{\hbar^2 \Gamma^2}{deg} \quad \text{c.g.s units}$$

[Aside: We will see that $\Gamma = \frac{4}{3} \left(\frac{\omega_{eg}}{C} \right)^3 \frac{deg^2}{\hbar}$
 $\Rightarrow deg^2 = \frac{3\hbar\Gamma}{4} \left(\frac{C}{\omega_{eg}} \right)^3$]

$$\Rightarrow I_{sat} = \left(\hbar \omega_{eg} \right) \left(\frac{1}{6\pi \left(\frac{C}{\omega_{eg}} \right)^2} \right) \left(\frac{\Gamma}{2} \right) = \frac{\text{Energy}}{\text{time} \cdot \text{Area}}$$

Here I have used $\sigma_{abs} = 6\pi \left(\frac{C}{\omega_{eg}} \right)^2$ = absorption cross-section

$$I_{sat} = \frac{\hbar \omega_{eg}}{\sigma_{abs}} \frac{\Gamma}{2}$$

for 2-level atom

Off-resonance

$$S = \frac{S_0}{1 + \frac{4\Delta^2}{\Gamma^2}} = \frac{I/I_{\text{sat}}}{1 + \frac{4\Delta^2}{\Gamma^2}}$$

Saturation falls off like $\frac{\Gamma^2}{\Delta^2}$ for $\Delta \gg \Gamma$

Absorption cross-section and scattering

Recall definition of cross-section:

Given incident intensity I , the absorbed power is

$$\begin{aligned} P_{\text{abs}} &= I \sigma_{\text{abs}} \\ &= (\text{Rate of absorption}) \times \hbar\omega \end{aligned}$$

$$\text{Rate of absorption} = \Gamma_{\text{stim}} = \frac{S}{2} \Gamma$$

$$\Rightarrow \frac{S\Gamma}{2} \hbar\omega = I \sigma_{\text{abs}} = \left(\frac{I/I_{\text{sat}}}{1 + \frac{4\Delta^2}{\Gamma^2}} \right) \frac{\Gamma}{2} \hbar\omega$$

$$\Rightarrow \text{On resonance} \quad I \sigma_{\text{abs}} = \left(\frac{I}{I_{\text{sat}}} \right) \frac{\Gamma}{2} \hbar\omega_{\text{eg}}$$

$$\Rightarrow \boxed{I_{\text{sat}} = \frac{\hbar\omega_{\text{eg}}}{\sigma_{\text{abs}}} \frac{\Gamma}{2}}$$

As before

For two level atom rate of absorption \rightarrow Photon is Scattered

$$\Rightarrow \text{Scattering rate} \quad \boxed{\frac{S\Gamma}{2} = \frac{I}{\hbar\omega} \sigma_{\text{abs}}}$$

Saturation and Power Broadening

Let us return to the steady-state coherence and polarizability. The lineshape we found was for weak excitation, i.e., $s \ll 1$

More generally

$$\rho_{ge}^{ss} = \left(\frac{+\sqrt{2}/2}{\Delta + i \Gamma/2} \right) \left(\rho_{e,e}^{ss} - \rho_{g,g}^{ss} \right)$$

\Rightarrow Steady state Bloch vector components

$$W^{ss.} = \rho_{e,e}^{ss.} - \rho_{g,g}^{ss.} = \begin{bmatrix} -1 \\ \frac{-1}{1+s} \end{bmatrix}$$

$$U^{ss.} = 2 \operatorname{Re}(\rho_{g,e}^{ss.}) = \left(\frac{\sqrt{2}\Delta}{\Delta^2 + \Gamma^2/4} \right) \left(\frac{1}{1+s} \right)$$

$$\Rightarrow U^{ss.} = -\frac{2\Delta}{\sqrt{2}} \frac{s}{1+s}$$

$$V^{ss.} = 2 \operatorname{Im}(\rho_{ge}^{ss.}) = \left(\frac{\sqrt{2}\Gamma/2}{\Delta^2 + \Gamma^2/4} \right) \left(\frac{1}{1+s} \right)$$

$$\Rightarrow V^{ss.} = -\frac{\Gamma}{\sqrt{2}} \frac{s}{1+s}$$

For $s \ll 1$

$$\begin{cases} U^{ss.} \approx -\frac{2\Delta}{\sqrt{2}} s = \frac{\Delta\sqrt{2}}{\Delta^2 + \frac{\Gamma^2}{4}} \\ -V^{ss.} \approx +\frac{\Gamma}{\sqrt{2}} s = \frac{+\Gamma\sqrt{2}/2}{\Delta^2 + \frac{\Gamma^2}{4}} \end{cases}$$

For $s \gg 1$, $\omega \rightarrow 0$ Saturation


$$\sqrt{2} \gg \Delta, \Gamma \Rightarrow u, v \rightarrow 0$$

\Rightarrow Saturation = Completely mixed state

More details:

$$\frac{s}{1+s} = \frac{\frac{I}{I_{\text{sat}}}}{1 + \frac{4\Delta^2}{\Gamma^2} + \frac{I}{I_{\text{sat}}}} = \frac{\frac{2\Omega^2}{\Gamma^2}}{1 + \frac{4\Delta^2}{\Gamma^2} + \frac{2\Omega^2}{\Gamma^2}}$$

$$\Rightarrow \frac{s}{1+s} = \frac{\sqrt{2}/2}{\Delta^2 + \frac{\Gamma^2}{4}(1 + \frac{I}{I_{\text{sat}}})}$$



"Power Broadened line width"

$$\boxed{\Gamma(I) = \Gamma \sqrt{1 + \frac{I}{I_{\text{sat}}}}}$$